

COLONOSCOPY

The purpose of the colonoscopy is to investigate the causes of disorders of the colon and rectum.

The procedure is carried out by passing a flexible endoscope through the anus and into the colon.

You will hear the results of the procedure in 4 to 6 weeks from the physician who referred you for it, unless otherwise specified.

Due to potential medication, we recommend that you do not drive to the procedure yourself.

Please cancel your appointment **at least one week before the procedure**, if you cannot attend at the scheduled time.

Before the procedure:

- Insulin-dependent diabetes: Please contact a diabetes nurse, if necessary.
- If you are on blood thinners, such as Marevan, Plavix, Pradaxa, Klexane, Fragmin, please contact the endoscopy unit or follow these instructions:

Stop taking the medication _____ INR _____ or according to separate instructions.

- Iron deficiency medication: stop taking for **one week**.
- Fibre products (such as Vi-Siblin, Agiocur): stop taking for **one week**.
- **Constipation:** take, for example, Movicol 2–3 sachets/day or Levolac 30 ml x1 in the evenings
3-5 days before starting cleansing. Available from pharmacies without prescription.

Diet one week before the procedure:

You are allowed to eat potatoes, rice, pasta, fish, chicken, meat, porridge and seed-free bread normally. You are allowed to eat fruit and vegetables as long as the seeds and skins are removed.

Herttoniemi Service Point

Siilikuja 3
00800 Helsinki
09 310 47692
Mon–Thu 8–14
Fri 8–13

**On the day of the procedure after office hours tel. 040
334 6960**

Malmi Service Point

Talvelantie 6, Malmi Hospital
00700 Helsinki
09 310 47692
Mon–Thu 8–14
Fri 8–13

**On the day of the procedure after office hours tel. 050
402 5789**

You are not allowed to eat: berries, nuts, seeds, grains, brans and similar foods.

EVENING INSTRUCTIONS

COLONSTERIL CLEANSING

For cleansing your colon, please purchase **six containers** of the prescription-free **Colonsteril** bowel cleansing product from your local pharmacy.

Please follow the instructions below for cleansing (instead of those provided with the product).

1. On the day before the procedure

- You may have breakfast and a light lunch normally, by no later than 14:00. During the afternoon and evening, you may freely consume liquid foods and clear liquids (such as water, juice, soft drinks, tea, coffee and meat, fish or vegetable broth) and take all your regular medication. **You must not eat any solid foods, only the clear liquids mentioned above are allowed.**

2. On the day of the procedure

- **Start drinking Colonsteril at 7:00.**
Mix one full container of the drops into one litre of water and drink it within one hour.
- During the day, drink a total of **6 litres of Colonsteril 3–4 hours before coming in for the procedure.**
- After drinking the cleansing product, it is important to drink the clear liquids listed above.
- You can make drinking the cleansing product easier by sucking on peppermint drops, for example.
- Take your regular medication, but do not take any tablet-based diabetes medication.

Herttoniemi Service Point

Siilikuja 3
00800 Helsinki
09 310 47692
Mon–Thu 8–14
Fri 8–13

**On the day of the procedure after office hours tel. 040
334 6960**

Malmi Service Point

Talvelantie 6, Malmi Hospital
00250 Helsinki
09 310 47692
Mon–Thu 8–14
Fri 8–13

**On the day of the procedure after office hours tel. 050
402 5789**