
COLONOSCOPY

The purpose of the colonoscopy is to investigate the causes of disorders of the colon and rectum.

The procedure is carried out by passing a flexible endoscope through the anus and into the colon. The mucous membranes of the colon are usually biopsied.

You will hear the results of the procedure in 4 to 6 weeks from the physician who referred you for it, unless otherwise specified.

Due to potential medication, we recommend that you do not drive to the procedure yourself.

Please cancel your appointment **at least one week before the procedure**, if you cannot attend at the scheduled time.

Before the procedure:

- Insulin-dependent diabetes: Please contact a diabetes nurse, if necessary.
- If you are on blood thinners, such as Marevan, Plavix, Pradaxa, Klexane, Fragmin, please contact the endoscopy unit or follow these instructions:

Stop taking the medication _____ INR _____ or according to separate instructions.

- Iron deficiency medication: stop taking for **one week**.
- Fibre products (such as Vi-Siblin, Agiocur): stop taking for **one week**.
- **Constipation:** if you suffer from constipation, take, for example, Movicol 2–3 sachets/day or Levolac 30 ml x1 in the evenings for 3–5 days before starting cleansing (available from pharmacies without prescription).

Diet one week before the procedure:

You are allowed to eat potatoes, rice, pasta, fish, chicken, meat, porridge and **seed-free** bread normally. You are allowed to eat fruit and vegetables as long as the **seeds and skins** are removed.

You are not allowed to eat: berries, nuts, seeds, grains, brans and similar foods.

Malmi Internal Medicine Outpatient Clinic
Siilikuja 3
00800 Helsinki
09 310 67269
Mon–Thu 8–11 and 12–14:30
Fri 8–11 and 12–13

Laakso Internal Medicine Outpatient Clinic
Lääkärintäti 8 L, 4th floor
00250 Helsinki
09 310 46194
Mon, Wed, Thu 8:15–11 and 12–14
Tue 9–11 and 12–14, Fri 8:15–11 and 12–13

PICOPREP CLEANSING

For cleansing your colon, please purchase one package of the prescription-free **Picoprep** bowel cleansing solution from your local pharmacy.

Two days before the procedure, eat only light meals, such as soup.

Please follow **the instructions below** for cleansing (instead of those provided with the product).

- **On the day before the procedure**

- **You must not eat any solid foods, only clear liquids** (meat, fish or vegetable broth; pulp-free juice; soft drinks; tea; coffee; **no** dairy products) and juice-based dietary supplements (such as Nutridrink) **are allowed**.
- **Drink the first Picoprep dose at 18:00.** Mix the Picoprep sachet into a glass of water. Mix the powder thoroughly for 2–3 minutes before drinking. After drinking, wait ½ an hour before drinking any other liquids in order to avoid diluting the cleansing product.
- After drinking the cleansing product, drink **at least 3 litres** of the aforementioned clear liquids and take your regular evening medication.

- **On the day of the procedure**

- **Continue the cleansing process by drinking the second Picoprep dose 4 hours before the procedure.**
- Wait ½ an hour. After this, drink **2 litres** of the aforementioned clear liquids at least 2 hours before coming in for the procedure.
- Take your necessary medication (such as blood pressure, asthma or epilepsy medication)
- **Do not take any tablet-based diabetes medication.**